

VANCOMYCIN PHARMACOKINETIC MONITORING IN CRITICALLY ILL NEONATAL PATIENTS

4CPS-161



D. Pascual Carbonell¹, J. Bodega Azuara¹, M. Martín Marqués¹, H. Suñer Barriga¹, I. Sacanella Anglès¹, C.D. Ciuci¹, P. López Broseta¹, A. García Molina¹, S. Conde Giner¹, I. Plo Seco¹, L. Canadell Vilarrasa¹


¹HOSPITAL UNIVERSITARIO JOAN XXIII, PHARMACY SERVICE, TARRAGONA, SPAIN

Salut/ HJ23
Hospital Joan XXIII

J01 – ANTIBACTERIALS FOR SYSTEMIC USE

BACKGROUND

VANCOMYCIN (glycopeptide)




Neonatal critically ill patients

Risk of infection
Dose adjustment

Pharmacokinetic monitoring



Neonatal critically ill patients with <1 month of life



AIM AND OBJECTIVES

Assessing possible under-exposure to vancomycin in critically ill neonatal patients after dosing, as recommended by standard guidelines.

- Retrospective observational

March 2021
June 2023



Tertiary hospital

Data collected from medical records

- Demographics
- Diagnosis
- Microbiological culture results
- Renal function
- Vancomycin dosing regimen
- Plasma concentration (PC)
- Antimicrobial treatment duration
- Occurrence of nephrotoxicity



Therapeutic PC VANCOMYCIN (before dose 4) 10-20 mg/dL

RESULTS



34 patients




79 pharmacokinetic determinations



45 treatments

Median duration of 6 days

31 (68,9%) empirical




- Catheter infection
- Sepsis
- Necrotising enterocolitis
- Suspected infection
- Pneumonia
- Septic shock
- Chorioamnionitis




28 positive microbiological cultures (62,2%)


Pathogens isolated in microbiological cultures



VANCOMYCIN PLASMA CONCENTRATION



Standard guidelines



48 under-treated (62,2%)

9 nephrotoxicity (13,3%)

CONCLUSIONS

It is necessary to review the recommended dosing regimens in this group of patients to achieve therapeutic PC of vancomycin from the start of treatment guided by pharmacokinetic monitoring.