

USE OF ERIBULIN IN LOCALLY ADVANCED OR METASTATIC BREAST CANCER

Contreras Rey MB, Garrido Martínez MT, Carrión Madroñal IM,
Yáñez Fera D, Sánchez Gómez E, Bocanegra Martín C

Juan Ramón Jiménez Hospital, Pharmacy. Huelva, Spain

4CPS-125

L01 - Cytostatics



BACKGROUND

Eribulin has been approved for locally advanced or metastatic breast cancer after at least one previous chemotherapy regimen for advanced disease, including an anthracycline and a taxane.

OBJECTIVES


To evaluate the effectiveness and safety of eribulin in a tertiary level hospital.

METHODS


- Retrospective observational study.
- Patients treated with eribulin (01 February 2014 - 11 October 2017).
- Variables: age, number of cycles, duration of treatment, number and type of previous chemotherapy regimens, progression-free survival (PFS), reported adverse events (AEs), dose reductions and dose delays between cycles.
- Data obtained from: electronic clinical records and the chemotherapy management software.

RESULTS

- 24 patients included, mean age 50.9 years (SD 9.4, range 32-67). At the data analysis, 4 were still in treatment with eribulin and 20 had finished it: median duration 3.15 months (4.5 cycles, range 1-8).
- Median of previous chemotherapy lines in locally advanced or metastatic stage: 3 (range 1-6).



Median PFS in the 17 patients who progressed during or after eribulin (but without having received a later treatment) was 2.8 months



- 62.5% of patients had any AE
- 1 patient interrupted the treatment due to AEs
- In patients who finished treatment, there were 2 delays because of neutropenia and 3 dose reductions due to toxicity

CONCLUSIONS

- In our patients, eribulin median PFS was lower than in EMBRACE trial. It could be explained because our patients received more previous regimens of chemotherapy for metastatic disease. In addition, our sample size was smaller.
- Regarding safety, eribulin was well tolerated and in most cases the AEs didn't forced to interrupt treatment.