



Hospital formulations for the treatment of nonalbicans vulvovaginitis

B. San José, Z. Baskaran, L. Serrano, I. Bilbao, B. Sordo, A. Bustinza, M. Castaño, B. Baza, S. Sautua, E. Rodriguez Hospital Universitario de Cruces, Barakaldo (Spain) begona.sanjoseruiz@osakidetza.net


Background


As vaginal flucytosine is not commercially available, it is necessary to prepare this formulation at Pharmacy Service.

Porpouse:

To describe the elaboration of flucytosine 15.5% (FLUCY15.5), and flucytosine 17% + amphotericin 3% (FLUAM) creams for vaginal use and their efficacy in the treatment of *Candida glabrata* and *Candida tropicalis*.

Materials and methods:





Conclusion

Local treatment with flucytosine and amphotericin was effective against vaginal infections caused by nonalbicans Candida species.

Disclosure: that may have

Authors of this presentation have the following to disclose concerning possible financial or personal relationship with comercial entities that may have a direct or indirect interest in the subject matter of this presentation:

8. San José Nothing to disclose 2. Baskaran: Nothing to disclose 3. Entiting to disclose 1. Elithor. Nothing to disclose 3. Elithor Nothing to disclose 3. Saturals Nothing to disclose 3.